


**Indian Journal of Chemistry**  
**Sect. A: Inorganic, Bio-inorganic, Physical, Theoretical & Analytical**  
[www.niscair.res.in](http://www.niscair.res.in); <http://nepr.niscair.res.in>  
 CODEN: ICACFC; ISSN: 0376-4719 (Print), 0975-0975 (Online)

**VOLUME 55A****NUMBER 6****JUNE 2016****CONTENTS**

- 649 Phosphorous-doped porous carbon derived from paste of newly growing *Ficus benghalensis* as hydrogen storage material**


Synthesis of heteroatom (P) doped porous carbon (PDPC) derived from the paste of newly growing Indian banyan tree (*Ficus benghalensis*), involving activation, carbonization and phosphorous-doping processes using  $H_3PO_4$  as activating agent and as phosphorous source, is reported. PDPC shows a wafer-like morphology with specific surface area of  $1406 \text{ m}^2/\text{g}$  and hydrogen storage capacity of ~1.2 wt% at 298 K and 100 bar.



Arjunan Ariharan, Balasubramanian Viswanathan\* &  
 Vaiyapuri Nandhakumar

- 657 Computational studies on chemically engineered carbon nanotubes as HCl sensor**


The effect of functionalization on sensing properties of SWNT on work function and HOMO-LUMO gap changes is studied by DFT calculations. Protonation of the basic group induces electron transfer from the semiconducting SWNTs, thereby altering the HOMO and LUMO levels and significantly changing the electronic properties and work function.



Taghi Haghtalab & Hamed Soleymanaabadi\*

**664 Temperature dependent study of thermophysical and optical properties of binary mixtures of imidazolium based ionic liquids**

The excess thermodynamic parameters and deviations in refractive index of  $[\text{Bmim}][\text{PF}_6]$ - $\alpha,\omega$ -propanediol binary mixtures indicate significant interactions in these mixtures, which follows the order: 1,2-propanedio > 1,3-propanediol. The results indicate the applicability of various mixing rules of refractive index in theoretically estimating refractive indices of these mixtures.




T S Krishna, M G Sankar, A K Nain\* &  
B Munibhadrayya

### Notes

**676 Sodium paramolybdate revisited**


A rational synthesis of  $\text{Na}_6[\text{Mo}_7\text{O}_{24}] \cdot 14\text{H}_2\text{O}$  and its spectral, thermal and electrochemical characterization is reported. Cyclic voltammetric and conductivity studies of an aqueous solution of  $\text{Na}_6[\text{Mo}_7\text{O}_{24}] \cdot 14\text{H}_2\text{O}$  reveal the presence of hydrated  $\text{Na}^+$  ions and heptamolybdate anions in solution, unlike a sodium coordinated heptamolybdate in the solid state.



Bilkhandarkoil R Srinivasan\* & Sudesh M Morajkar

**681 Phenoxazinone synthase activity of a mononuclear cobalt(III) complex**


The trivalent cobalt complex,  $[\text{Co}(\text{HL})_2](\text{OAc})\cdot\text{H}_2\text{O}$  (**1**) [ $\text{H}_2\text{L} = N-(2\text{-hydroxyethyl})-3\text{-methoxysalicylaldimine}$ ], is an effective model for phenoxazinone synthase to oxidize  $\alpha$ -aminophenol to 2-aminophenoxyazine-3-one in methanol in presence of aerial oxygen. The reaction follows Michaelis-Menten enzymatic reaction kinetics with high turnover number ( $K_{cat} = 2.83 \times 10^5 \text{ h}^{-1}$ ).



Merry Mitra & Rajarshi Ghosh\*

**686 One-step synthesis of ZnO nanowires on zinc foils and their photocatalytic properties**


ZnO nanostructures with different morphologies have been grown on zinc foil substrate by a novel and facile hydrothermal method, without the assistance of any catalyst or template. The resulting ZnO nanorods have diameter of ~70–90 nm and average length in the range of 0.5–2.0  $\mu\text{m}$ . The as-grown ZnO samples have very good crystallinity. A comparison of the photocatalytic degradation of methyl orange with different ZnO shows that the photocatalytic properties of the ZnO nanostructures depend on the morphology of ZnO. The growth process of the ZnO nanorods is proposed based on the solid-liquid-solid mechanism.



Mohamad Mohsen Momeni\*

- 692 Naphthalimide based hybrid nanoparticles for selective recognition of Hg(II)

Anu Saini & Navneet Kaur\*



- 698 Corrigendum

Authors for correspondence are indicated by (\*).

## Online Submission for IJCA Articles

Articles submitted for consideration of publication in the Indian J Chem, Sec A, should be uploaded through our Online Submission portal, <http://op.niscair.res.in>. Articles submitted by email or postal services shall not be processed.

Authors are requested to register at <http://op.niscair.res.in>. The articles (in MS Word) appended with the certificate of originality, illustrations (jpeg/tiff) and supplementary data, if any, may be uploaded through the User Home page. Further information is available at [http://www.niscair.res.in/jinfo/author\\_guidelines\\_online\\_submission.pdf](http://www.niscair.res.in/jinfo/author_guidelines_online_submission.pdf).

For technical support and assistance for online submission, Shri Sanjay Burde may be contacted ([sanjayburde@niscair.res.in](mailto:sanjayburde@niscair.res.in)); Tel.: 91-11-25846301.